


AMENDMENTS TO THE CLAIMS


This following listing of claims will replace all prior listings for the application.

Listing of claims:


1. (currently amended) A process for the production of an optically active amino alcohol represented by the following formula (I)



(wherein, R^2 , R^3 , R^4 and R^5 each independently is a hydrogen atom, a lower alkyl group or an optionally-substituted phenyl group; R^2 or R^3 may be bonded to R^4 or R^5 forming a ring together with the adjacent carbon atoms; A^1 is $-(CH_2)_m-$ while A^2 is $-(CH_2)_n-$ (where m and n each is an integer of 0 to 3 and $m + n$ is 1 to 3); * is an asymmetric carbon atom where the relative configuration of hydroxyl group to ~~alkoxycarbonyl~~ amino group on each of the asymmetric carbons marked * is trans) or a salt thereof, comprising reacting an optically active hydroxycarboxylate represented by the following formula (IV)




(wherein, R^1 is an alkyl group having 1 to 6 carbon(s); R^2 , R^3 , R^4 , R^5 , A^1 , A^2 , m , n and * have the same meanings as defined above where the relative configuration of hydroxyl group to ~~amino~~ alkoxycarbonyl group on each of the asymmetric carbons marked * is trans) with hydrazine to prepare an optically-active hydroxycarboxylic hydrazide compound represented by the following formula (III)



(wherein, R^2 , R^3 , R^4 , R^5 , A^1 , A^2 , m , n and $*$ have the same meanings as defined above where the relative configuration of hydroxyl group to hydrazinocarbonyl group on each of the asymmetric carbons marked $*$ is trans), then conducting a Curtius reaction in the presence of an alcohol represented by the following formula (VI)




(wherein, R^6 is an alkyl group having 1 to 6 carbon(s) or an optionally-substituted benzyl group) to give an optically active alkoxy carbonylamino alcohol represented by the following formula (II)




(wherein, R^2 , R^3 , R^4 , R^5 , R^6 , A^1 , A^2 , m , n and $*$ have the same meanings as defined above where the relative configuration of hydroxyl group to alkoxy carbonylamino group on each of the asymmetric carbons marked $*$ is trans) and then deprotecting a protective group for the amino group.


2. (currently amended) A process for the production of an optically active alkoxy carbonylamino alcohol represented by the following formula (II)



(wherein, R^2 , R^3 , R^4 and R^5 each independently is a hydrogen atom, a lower alkyl group or an optionally-substituted phenyl group; R^2 or R^3 may be bonded to R^4 or R^5 forming a ring together with the adjacent carbon atoms; R^6 is an alkyl group having 1 to 6 carbon(s) or an optionally-substituted benzyl group; A^1 is $-(CH_2)_m$ - while A^2 is $-(CH_2)_n$ - (where m and n each is an integer of 0 to 3 and $m + n$ is 1 to 3); * is an asymmetric carbon atom where the relative configuration of hydroxyl group to alkoxy carbonyl group on each of asymmetric carbons marked * is trans), comprising reacting an optically active hydroxycarboxylate represented by the following formula (IV)



(wherein, R^1 is an alkyl group having 1 to 6 carbon(s); R^2 , R^3 , R^4 , R^5 , A^1 , A^2 , m , n and * have the same meanings as defined above where the relative configuration of hydroxyl group to alkoxy carbonyl group on each of the asymmetric carbons marked* is trans) with hydrazine to prepare an optically-active hydroxycarboxylic hydrazide compound represented by the following formula (III)




(wherein, R^2 , R^3 , R^4 , R^5 , A^1 , A^2 , m , n and $*$ have the same meanings as defined above where the relative configuration of hydroxyl group to hydrazinocarbonyl group on each of the asymmetric carbons marked $*$ is trans) and conducting a Curtius reaction in the presence of an alcohol represented by the following formula (VI)




(wherein, R^6 is an alkyl group having 1 to 6 carbon(s) or an optionally-substituted benzyl group).

3. (previously presented) The process according to claim 1 or 2, wherein the optically active hydroxycarboxylate represented by the following formula (IV)



(wherein, R^1 , R^2 , R^3 , R^4 , R^5 , A^1 , A^2 , m , n and $*$ have the same meanings as defined above where the relative configuration of hydroxyl group to alkoxycarbonyl group on each of the asymmetric carbons marked $*$ is trans) is a product prepared by subjecting a β -keto ester represented by the following formula (V)



(wherein, R^1 , R^2 , R^3 , R^4 , R^5 , A^1 , A^2 , m and n have the same meanings as defined above) to an asymmetric hydrogenation in the presence of a ruthenium complex including an optically active phosphine compound as a ligand.

4. (previously presented) The process according to claim 1 or 2, wherein R^6 is an optionally substituted benzyl group.

5. (previously presented) The process according to claim 1 or 2, wherein R⁶ is a benzyl group.
6. (previously presented) The process of claim 3 wherein R⁶ is an optionally substituted benzyl group.
7. (previously presented) The process of claim 3 wherein R⁶ is a benzyl group.